
CORRELATIONS BETWEEN METTABILITY AND STRUCTURE IN MONOLAYERS OF ALKANETHIOLS ADSORBED ON GOLD(U) HARVARD UNIV CAMBRIDGE MASS DEPT OF CHEMISTRY C D BAIN ET ALMAY 88 TR-7 N00014-85-K-0898 F/G 7/4 1/1 AD-R194 348 UNCLASSIFIED NL



AD-A194 348

CORRELATIONS BETWEEN WETTABILITY AND STRUCTURE IN MONOLAYERS OF ALKANETHIOLS ADSORBED ON GOLD

Colin D. Bain and George M. Whitesides*
Department of Chemistry
Harvard University
Cambridge MA 02138

Technical Report No. 7 (May 1988)

Interim Technical Report

(Accepted for publication in <u>J. Am. Chem. Soc.</u>)

PREPARED FOR DEFENSE ADVANCED RESEARCH PROJECTS AGENCY 1400 Wilson Boulevard Arlington VA 22209

DEPARTMENT OF THE NAVY
Office of Naval Research, Code 1130P
800 North Wuincy Street
Arlington VA 22217-5000

ARPA Order No.: NR 356-856

Contract No.: N00014-85-K-0898 Effective Date: 85 September 01 Expiration Date: 88 August 31

Principal Investigator: George M. Whitesides

(617) 495-9430

The views and conclusions in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U.S. Government.

Approved for public succession Distribution I elimited



REPRODUCED AT GOVERNMENT EXPENSE

| SECURITY CLASSIFICATION OF THIS PAGE | | | | | |
|---|--|---|--|---|------------------------------------|
| | REPORT DOCU | MENTATION | PAGE | | |
| Ta REPORT SECURITY CLASSIFICATION Unclassified | | 16 RESTRICTIVE MARKINGS | | | |
| 2a SECURITY CLASSIFICATION AUTHORITY | | 3 DISTRIBUTION, AVAILABILITY OF REPORT Approved for public release; distributio unlimited | | | |
| 26 DECLASS:F CATION / DOWNGRADING SCHEDULE | | | | | |
| 4 PERFORMING ORGANIZATION REPORT NUMBER(S) | | 5 MONITORING ORGANIZATION REPORT NUMBER(S) | | | |
| Technical Report #7 | | 1 | | | |
| 6a NAME OF PERFORMING ORGANIZATION 6b OFFICE SYMBOL | | 7a NAME OF MONITORING ORGANIZATION | | | |
| Harvard University | (If applicable) | Office of Naval Research | | | |
| 6c. ADDRESS (City, State, and ZIP Code) | 1. | 7b ADDRESS (City, State, and ZIP Code) | | | |
| Office of Sponsored Research Holvoke Center, Fourth Floor | | Code 11130P 800 North Quincy Street | | | |
| Cambridge MA 02138-4993 | | Arlington | | -5000 | |
| Ba NAME OF FUNDING/SPONSORING ORGANIZATION | 8b OFFICE SYMBOL (If applicable) | 9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER | | ION NUMBER | |
| ONR/DARPA 8c ADDRESS (City, State, and ZIP Code) | | 10 SOURCE OF | FUNDING NUMBE | 90 | |
| 800 North Quincy Street | | PROGRAM | PROJECT | TASK | WORK UN |
| Arlington VA 22217-5000 | | 85-K-0898 | NO. NR 356-856 | NO | ACCESSION |
| "Correlations between Wetta on Gold" 12 PERSONAL AUTHOR(S) Colin D. B | ain and George M. | | ayers or Aik | | |
| 13a TYPE OF REPORT 13b TIM Interim FROM | E COVERED TO | 14 DATE OF REPORT (Year, Month, Day) 15. PAGE 6 May 1988 | | PAGE COUNT | |
| 16 SUPPLEMENTARY NOTATION | | | | | |
| | | (Continue on reverse if necessary and identify by block number) | | | |
| F-ELD GROUP SUB-GROUP | monolayer gold | weecability sell assembly | | | bly XPS |
| 19 ABSTRACT (Continue on reverse if necess | | | | 1 | |
| Long-chain ak monolayers. Coads solutions of varyi of which is correl Adsorption of the and formation of m favored over mixed | ng composition gē ated closely with longer chain thio onolayers compris |) 10H and HS nerates mono the composi 1 is preferr ing predomin | (CH' ₂) 21 CH ₃ flayers, the tion of the ed over the antly one co | rom eth wettabi monolay short component | nanolic ility ver. chain, |
| 20 DISTRIBUTION AVAILABILITY OF ABSTRA ☑ UNCLASS FIED UNLIMITED ☐ SAME A | | | ECURITY CLASSIFI | CATION | |
| 22a GAME OF RESPONSIBLE NOIVIDUAL Kenneth J. Wynne | <u> </u> | | (Include Area Cod | (e) 22c O | FFICE SYMBOL |

DD FORM 1473, 84 MAR

Correlations between Wettability and Structure in Monolayers of Alkanethiols Adsorbed on Gold¹

Colin D. Bain² and George M. Whitesides*

Department of Chemistry.

Harvard University.

Cambridge, MA 02138

Abstract

Long-chain alkanethiols adsorb onto gold from solution and form monolayers. Coadsorption of HS(CH₂)₁₁OH and HS(CH₂)₂₁CH₃ from ethanolic solutions of varying composition generates monolayers, the wettability of which is correlated closely with the composition of the monolayer. Adsorption of the longer-chain thiol is preferred over the short chain, and formation of monolayers comprising predominantly one component is favored over mixed monolayers containing both component thiols.



| Access | ion For | r | _/ | |
|--|---------|------------|------|---|
| NTIS | T.A&I | | ZZ. | |
| DIIC I | AB | | | |
| Unanio | | | | |
| Justia | 105110 | n | | |
| | | | | |
| By | | | | _ |
| Distri | bution | / | | |
| Avai | abilit | y C | ode5 | |
| - | Avall | and/ | 'or | |
| Dist | Spec | ial | | |
| 1 | | 1 | | |
| | | 1 | | |
| IN | | l | | |
| 1 | <u></u> | ┸—— | | |

Long-chain thiols (HS(CH₂)_nX) adsorb from solution onto gold surfaces and form well-packed, ordered, oriented monolayers.^{3,4} The sulfur coordinates strongly to the gold, the polymethylene chains are all-trans and tilted ~20–30° from the normal to the surface,^{4,5} and the tail group, X, is the predominant group exposed at the monolayer/liquid or monolayer/air interface.⁶ Coadsorption of two or more thiols differing in tail group or chair length provides a flexible system for varying the chemistry and structure of the surface in a controlled and pre-determined way.⁷ In this paper we use monolayers comprising a mixture of HS(CH₂)₁₁OH (represented as HSC₁₀CH₂OH to emphasize the two important variables: chain length and tail group) and HS(CH₂)₂₁CH₃ (HSC₂₁CH₃) on gold to demonstrate a relationship between the microscopic structure of the surface and the wettability of the monolayer (Figure 1). This work is part of a program of physical-organic chemistry designed to relate atomic-level structure of a surface to macroscopic physical properties such as wetting and adhesion.⁸

Monolayers were formed by immersing gold-mirror substrates (prepared by evaporation of gold onto chromium-primed, polished silicon wafers) in solutions of the thiols in degassed ethanol for 12 hours⁹ at room temperature.⁶ The composition of the monolayer was controlled by varying the ratio $R = [HSC_{10}CH_2OH]/[HSC_{21}CH_3]$ in solution, with the total concentration of thiols held constant at 1 mM. We used two independent techniques, optical ellipsometry and X-ray photoelectron spectroscopy (XPS), to measure the composition of the monolayer. Since the two thiols differ in chain length, the relative thickness obtained from ellipsometry and from the peak area of gold¹⁰ in XPS reflects this composition. Similarly, the difference in tail groups allows us to calculate the surface concentration of $HSC_{10}CH_2OH$ from the peak area of oxygen. The advancing contact angles (θ_a) of water and hexadecane (HD) provide useful measures of wettability. Since the pure methyl-terminated monolayer is hydrophobic ($\theta_a(H_2O) = \theta_a(HD) = 0^\circ$), and the alcohol-terminated monolayer is hydrophilic ($\theta_a(H_2O) = \theta_a(HD) = 0^\circ$),

contact angles are very sensitive to the composition of the monolayer and the structure of the surface.

Figure 2 plots the ellipsometric thickness, contact angles of water and hexadecane, and XPS peak areas of gold and oxygen against R.¹¹ Here we note two salient features of these graphs: a detailed discussion will be deferred to a subsequent paper. First, ellipsometry and XPS peak areas indicate a dramatic change in the composition of the monolayer over a narrow range of solution composition, R = 7 - 20. This change in the composition of the monolayer is closely correlated with a sharp increase in the hydrophilicity and oleophilicity of the surface as measured by the advancing contact angles of water and hexadecane, respectively.¹² Thus the structure of the surface on a microscopic scale is clearly and directly linked to the wettability of the monolayer, an important macroscopic quantity. Second, the inflection in the curves occurs, not at R = 1, but at $R \approx 11.^{13,14}$ This difference between solution and surface compositions is a general feature of competitive adsorption experiments.¹⁵ In this experiment, Van der Waals forces between close-packed hydrocarbon chains favour adsorption of the longer-chain thiol.

We attempted to model the composition of the monolayer by a simple equilibrium expression (Eq. 1) between the solution and the surface (dotted curve in Figure 2, upper graph)

$$HSC_{10}CH_2OH_{surface} + HSC_{21}CH_{3solution} \rightleftharpoons HSC_{10}CH_2OH_{solution} + HSC_{21}CH_{3surface}$$
 (1)

where the equilibrium constant $K_{eq} = 11$, independent of solution concentration. Clearly the observed data do not follow this simple expression: the two components of the monolayer do not act independently, and may act cooperatively to minimize the free energy. Monolayers composed predominantly of the long-chain methyl-terminated thiol $HSC_{21}CH_3$ (maximizing the chain-packing interactions) or the short-chain hydroxy-terminated thiol $HSC_{10}CH_2OH$ (maximizing H-bonding both with the solvent and within the monolayer)

are preferred over monolayers containing a mixture of the thiols.¹⁶ The data do not, however, take the form of step functions in R, which would be expected thermodynamically if the formation of macroscopic islands were favored.¹⁷ Over a narrow range of R it is possible to form intermediate monolayers containing both thiols; the exact structure of these monolayers is not clear, but the data are consistent with a model in which the two components segregate into small clusters on the surface. These monolayers provide a model system for studying the wettability of complex structures, and for examining how polar groups interact to minimize their energy in a non-polar environment.¹⁸

Mixed monolayers of thiols on gold allow us to engineer ordered, two-dimensional systems with Å--level control over thickness and structure, and with chemical control over wettability. Synthetic variation of the tail groups and chain lengths provides great flexibility in the design of the interfacial structures and gives these systems wide applicability in the physical, biological and medical sciences.

Acknowledgement. We are grateful to R. Nuzzo and M. Wrighton for helpful discussions.

References and Notes

¹ Supported in part by the Office of Naval Research and the Defense Advanced Projects Research Agency.

XPS spectra were obtained using facilities obtained through the DARPA/URI and maintained in the Harvard University Materials Research Laboratory.

² IBM Pre-Doctoral Fellow in Physical Chemistry 1985-86.

³ Nuzzo, R. G.; Allara D. L. J. Am. Chem. Soc. 1983, 105, 4481-4483; Strong, L.; Whitesides, G. M. Langmuir in press.

⁴ Porter, M. D.; Bright, T. B.; Allara, D. L.; Chidsey, C. E. D. J. Am. Chem. Soc. 1987, 109, 3559-3568.

⁵ Nuzzo, R. G.; Dubois, L. H.; Allara, D. L. unpublished results.

⁶ Bain, C. D.; Troughton, E. B.; Tao, Y.-T.; Evall, J.; Whitesides, G. M.; Nuzzo, R. G. manuscript in preparation.

⁷ Bain, C. D.; Whitesides, G. M. Science in press.

⁸ Troughton, E. B.; Bain, C. D.; Whitesides, G. M.; Nuzzo, R. G.; Allara, D. L.; Porter, M. D. Langmuir 1988, 4, 000-000; Holmes-Farley, S. R.; Reamey, R. H.; McCarthy, T. J.; Deutch, J.; Whitesides, G. M. Langmuir 1985, 1, 725-40; Holmes-Farley, S. R.; Whitesides, G. M. Langmuir 1986, 2, 266-281; Holmes-Farley, S. R.; Whitesides, G. M. Langmuir 1987, 3, 62-75; Holmes-Farley, S. R.; Reamey, R. H.; Nuzzo, R. G.; McCarthy, T. J.; Whitesides, G. M. Langmuir 1987, 3, 799-815; Holmes-Farley, S. R.; Bain, C. D.; Whitesides, G. M. Langmuir in press.

⁹ For pure thiols, initial monolayer formation is very rapid (~few seconds) with limiting properties reached after a few hours.


¹⁰ The intensity of the photoelectron peak decreases exponentially with the thickness of the monolayer due to inelastic scattering of the photoelectrons (Briggs, D.; Seah, M. P *Practical Surface Analysis*; Wiley: Chichester, 1983).


- 11 We prefer logR to the mole fraction, χ , as the abscissa because kTlogR is related to the Gibbs free energy and hence this choice of axis highlights thermodynamic contributions to the adsorption process.
- ¹² Hexadecane and water interact predominantly by dispersion and polar forces, respectively. The detailed inferences from the contact angle data and comparisons with other monolayer systems will be discussed in a subsequent paper.
- ¹³ The ellipsometric thickness yields a value of $R(\chi_{1/2}) = 11$ where $\chi_{1/2} = 1:1$ ratio of the two thiols in the monolayer. XPS peak areas yield $R(\chi_{1/2}) = 10-12$ depending on the model used to analyze the data.
- ¹⁴ The monolayer system had not quite reached equilibrium when these measurements were made. Over a period of two weeks $R(\chi_{1/2})$ slowly increased to $R \approx 14$.
- 15 Bain, C. D.; Whitesides, G. M. unpublished results.
- Despite being enthalpically disfavored, some HSC₂₁CH₃ is incorporated into the HSC₁₀CH₂OH monolayer even at large R (and vice versa) due to the favorable entropy of mixing.
- ¹⁷ We believe that adsorption of thiols onto gold is controlled to a large extent by thermodynamics, although kinetics also play a role, as evinced by the slow change in the monolayer composition with time after the initial, rapid adsorption.
- 18 Israelachvili, J. N. Intermolecular and Surface Forces; Academic: New York, 1985.

FIGURE CAPTIONS

Figure 1. Schematic illustrations of monolayer structures: pure HS(CH₂)₁₁OH (A); pure HS(CH₂)₂₁CH₃ (B); monolayer containing a mixture of the two thiols (C).

Figure 2. Monolayers formed by the adsorption of mixtures of $HS(CH_2)_{11}OH$ and $HS(CH_2)_{21}CH_3$ onto gold from solution. The abscissa represents the ratio, $R = [HS(CH_2)_{11}OH]/[HS(CH_2)_{21}CH_3]$ in *solution*. Squares and circles represent data from two separate experiments. Upper Figure: ellipsometric thickness. The solid curve is a fit to the data. The dotted curve represents the theoretical thicknesses for $K_{eq} = 11$ (see text for definition). Middle Figure: advancing contact angles of water (open symbols) and hexadecane (HD) (solid symbols) obtained by the sessile drop technique. Lower Figure: areas of the Au $4f^7/_2$ (open symbols) and O 1s peaks (solid symbols) obtained by XPS. The vertical scale is arbitrary. Data were collected on a SSX-100 X-ray photoelectron spectrometer (Surface Science Instruments) with a monochromatized Al $K\alpha$ source, 100 eV pass energy, and 1-mm X-ray spot. The peaks were fitted using a symmetrical 90% Gaussian/ 10% Lorentzian profile.





CONTRACT NUMBER: NCCO14-85-K-08

CONTRACT DATA REQUIREMENTS LIST ATTACHMENT NO.: 1 INSTRUCTIONS FOR DISTRIBUTION ARPA/CNR

MINIMUM DISTRIBUTION OF TECHNICAL REPORTS

| | | NUMBER OF COPIES | | | |
|-------------------------------|--------|--|--|--|--|
| 227222 | ೨೦೨೩೩೨ | • | CNCLASSIFIED/LIMITED | | |
| • | CCCE | UNCLASSIFIED/UNLIMITED | AND CLASSIFIED | | |
| Director, Advanced Research | EX1241 | 2 | . 2 | | |
| Projects Agency | | | . | | |
| 1400 Wilson Boulevard | | | <i>₩</i> | | |
| Amilington, Vinginia 20009 | | | Mark Comment | | |
| ATTN: Program Management | | • | J. J. J. Co. | | |
| Scientific Officer | 41000K | · 3 | terre parious later de la later de later de later de la later de l | | |
| Administrative Contracting | 366016 | 1 | in in the second | | |
| Cfficer | | · | St. St. | | |
| Olrector, Naval Research | NC0173 | 5 g ^x | <u>.</u> | | |
| Laboratory, ATTN: Code 2627 | | | • | | |
| Washington, D. C. 20375 | | A Control of the Cont | | | |
| Defense Technical Information | S47031 | 12 | 2 | | |
| Canter | | | | | |
| Bldg. E, Cameron Station | | | | | |
| Alexandria, Virginia 22314 | | • | | | |

One (1) copy of each technical report resulting from work performed in the area of tactical technology shall be sent to:

TACTEC ECOD CAACOC Battelle Memorial Institue 505 King Avanue Columbus, Chio 43201

MINIMUM DISTRIBUTION OF REPORTS WHICH ARE NOT TECHNICAL REPORTS

| | | MUMPER OF C | CPIES |
|--|-----------------------|------------------------|--|
| ADDRESSES | DODAAD <u>CCDE</u> | UNCLASSIFIED/UNLIKITED | UNCLASSIFIED/LIMITED AND CLASSIFIED |
| Director, Advanced Research Projects Agency 1400 Wilson Boulevard Arlington, Virginia 22209 ATTN: Program Management | HX1241 | 2 | 2 |
| Scientific Officer | N00014 | 3 | 3 |
| Administrative Contracting | N66016 | 1 | 1 |

If the Scientific Cificer directs, the Contractor shall make additional distribute of technical reports and such other reports as may be specified by the Scientific Cificar in accordance with a supplemental distribution list provided by the Stiant Cfficer.

ABSTRACTS DISTRIBUTION LIST, 356B

Professor T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Kurt Baum Fluorochem, Inc. 680 S. Ayon Avenue Azuza, California 91702

Dr. Ulrich W. Suter Department of Chemical and Engineering Massachusetts Institute of Technologies Room E19-628 Cambridge, MA 02139-4309

Dr. William Bailey
Department of Chemistry
University of Maryland
College Park, Maryland 20742

Dr. J.C.H. Chien
Department of Polymer Science and
Engineering
University of Massachusetts
Amherst, MA 01003

Professor G. Whitesides Department of Chemistry Harvard University Cambridge, Massachusetts 02138

Dr. K. Paciorek Ultrasystems, Inc. P.O. Box 19605 Irvine, California 92715

Dr. Ronald Archer Department of Chemistry University of Massachusetts Amherst, Massachusetts 01002

Professor D. Seyferth
Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Professor J. Moore Department of Chemistry Rensselaer Polytechnic Institute Troy. New York 12181

Dr. V. Percec
Department of Macromolecular
Science
Case Western Reserve University
Cleveland, Ohio 44106

Dr. Gregory Girolami Department of Chemistry University of Illinois Urbana-Champagne, IL 61801

Dr. Ted Walton Chemistry Division Code 6120 Naval Research Lab Washington D.C. 20375-5000

Professor Warren T. Ford Department of Chemistry Oklahoma State University Stillwater, OK 74078

Professor H. K. Hall, Jr. Department of Chemistry The University Arizona Tucson, Arizona 85721

Dr. Fred Wudl Department of Chemistry University of California Santa Barbara, CA 93106

Professor Kris Matjaszewski Department of Chemistry Carnegie-Mellon University 4400 Fifth Avenue Pittsburgh, PA 15213

Professor Richard Schrock Department of Chemistry Massachusetts Institute of Technology Cambridge, MA 02139

ABSTRACTS DISTRIBUTION LIST, 356B

Professor A. G. MacDiarmid
Department of Chemistry
University of Pennsylvania
Philadelphia, Pennsylvania 19174

Dr. E. Fischer, Code 2853 Naval Ship Research and Development Center Annapolis, Maryland 21402

Professor H. Allcock
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Professor R. Lenz Department of Chemistry University of Massachusetts Amherst, Massachusetts 01002

Professor G. Wnek
Department of Chemistry
Rensselaer Polytechnic Institute
Troy, NY 12181

Professor C. Allen
Department of Chemistry
University of Vermont
Burlington, Vermont 05401

Dr. Ivan Caplan DTNSRDC Code 0125 Annapolis, MD 21401

Dr. R. Miller Almaden Research Center 650 Harry Road K91B801 San Jose, CA 95120

Dr. William B. Moniz Chemistry Division Naval Research Laboratory Washington, D.C. 20375-5000 Dr. Richard M. Laine SRI International 333 Ravenswood Avenue Menlo Park, California 94025

Dr. L. Buckley Naval Air Development Center Code 6063 Warminster, Pennsylvania 18974

Dr. James McGrath Department of Chemistry Virginia Polytechnic Institute Blacksburg, Virginia 24061

Dr. Geoffrey Lindsay Chemistry Division Naval Weapons Center China Lake, California 93555

Professor J. Salamone
Department of Chemistry
University of Lowell
Lowell, Massachusetts 01854

Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D. C. 20375-5000

Professor T. Katz Department of Chemistry Columbia University New York, New York 10027

Dr. Christopher K. Ober
Department of Materials Science
and Engineering
Cornell University
Ithaca, New York 14853-1501